Contents:

Volume 152, Number 1, January 8, 2001

In Brief

Wnt inhibits apoptosis.
Esp1p triggers both phases of anaphase.
Calreticulin and a termination factor in nuclear export.
Proteomics for phagosomes.
W.A. Wells

Mini-Review

F1 **Molecular machines: Putting the pieces together.**
E. Nogales and N. Grigorieff

Regular Articles

1 ***Wasp*, the *Drosophila* Wiskott-Aldrich syndrome gene homologue, is required for cell fate decisions mediated by *Notch* signaling.**
S. Ben-Yaacov, R. Le Borgne, I. Abramson, F. Schweisguth, and E.D. Schejter

15 **Replication origins in *Xenopus* egg extract are 5–15 kilobases apart and are activated in clusters that fire at different times.**
J.J. Blow, P.J. Gillespie, D. Francis, and D.A. Jackson

27 **A novel role of the budding yeast separin Esp1 in anaphase spindle elongation: Evidence that proper spindle association of Esp1 is regulated by Pds1.**
S. Jensen, M. Segal, D.J. Clarke, and S.I. Reed

41 **Vacuolar storage proteins are sorted in the cis-cisternae of the pea cotyledon Golgi apparatus.**
S. Hillmer, A. Movafeghi, D.G. Robinson, and G. Hinz

51 **Membrane recruitment of Aut7p in the autophagy and cytoplasm to vacuole targeting pathways requires Aut1p, Aut2p, and the autophagy conjugation complex.**
J. Kim, W.-P. Huang, and D.J. Klionsky

65 **Integrin activation by regulated dimerization and oligomerization of platelet endothelial cell adhesion molecule (PECAM)-1 from within the cell.**
T. Zhao and P.J. Newman

75 **A functional interaction between the survival motor neuron complex and RNA polymerase II.**
L. Pellizzoni, B. Charroux, J. Rappsilber, M. Mann, and G. Dreyfuss

87 **Wnt-1 signaling inhibits apoptosis by activating β-catenin/T cell factor–mediated transcription.**
S. Chen, D.C. Guttridge, Z. You, Z. Zhang, A. Fribley, M.W. Mayo, J. Kitajewski, and C.-Y. Wang

97 **Targeting Pyk2 to β1-integrin–containing focal contacts rescues fibronectin-stimulated signaling and haptotactic motility defects of focal adhesion kinase–null cells.**
C.K. Klingbeil, C.R. Hauck, D.A. Hsia, K.C. Jones, S.R. Reider, and D.D. Schlaepfer

111 **Differential localization of Rho GTPases in live cells: Regulation by hypervariable regions and RhoGDI binding.**[o]
D. Michaelson, J. Silletti, G. Murphy, P. D'Eustachio, M. Rush, and M.R. Philips

127 **Calreticulin is a receptor for nuclear export.**
J.M. Holaska, B.E. Black, D.C. Love, J.A. Hanover, J. Leszyk, and B.M. Paschal

141 **NXT1 is necessary for the terminal step of Crm1-mediated nuclear export.**
B.E. Black, J.M. Holaska, L. Lévesque, B. Ossareh-Nazari, C. Gwizdek, C. Dargemont, and B.M. Paschal

157 **The telomerase/vault-associated protein TEP1 is required for vault RNA stability and its association with the vault particle.**
V.A. Kickhoefer, Y. Liu, L.B. Kong, B.E. Snow, P.L. Stewart, L. Harrington, and L.H. Rome

165 **The phagosome proteome: Insight into phagosome functions.**
J. Garin, R. Diez, S. Kieffer, J.-F. Dermine, S. Duclos, E. Gagnon, R. Sadoul, C. Rondeau, and M. Desjardins

181 **Cbl associates with Pyk2 and Src to regulate Src kinase activity, $\alpha_v\beta_3$ integrin-mediated signaling, cell adhesion, and osteoclast motility.**
A. Sanjay, A. Houghton, L. Neff, E. DiDomenico, C. Bardelay, E. Antoine, J. Levy, J. Gailit, D. Bowtell, W.C. Horne, and R. Baron

197 **Mitotic spindle integrity and kinetochore function linked by the Duo1p/Dam1p complex.**
I.M. Cheeseman, M. Enquist-Newman, T. Müller-Reichert, D.G. Drubin, and G. Barnes

213 **The Sar1 GTPase coordinates biosynthetic cargo selection with endoplasmic reticulum export site assembly.**[o]
M. Aridor, K.N. Fish, S. Bannykh, J. Weissman, T.H. Roberts, J. Lippincott-Schwartz, and W.E. Balch

Report

231 **TANGLED1: A microtubule binding protein required for the spatial control of cytokinesis in maize.**[o]
L.G. Smith, S.M. Gerttula, S. Han, and J. Levy

[o]The online version of this article contains supplemental material.

Volume 152, Number 2, January 22, 2001

In Brief

Maintenance of mitochondrial DNA.
Timing of DNA replication.
Conserved mitotic spindle components identified.
Arterial injection of muscle-derived stem cells.
Voltage-dependent anion channel in apoptosis.

A.W. Dove

Comment

F11 **Nuclear position leaves its mark on replication timing.**

D.M. Gilbert

Regular Articles

237 **Essential role of voltage-dependent anion channel in various forms of apoptosis in mammalian cells.**

S. Shimizu, Y. Matsuoka, Y. Shinohara, Y. Yoneda, and Y. Tsujimoto

251 **Cortical Num1p interacts with the dynein intermediate chain Pac11p and cytoplasmic microtubules in budding yeast.**

M. Farkasovsky and H. Küntzel

263 **Activation of endogenous thrombin receptors causes clustering and sensitization of epidermal growth factor receptors of Swiss 3T3 cells without transactivation.**

M.F. Crouch, D.A. Davy, F.S. Willard, and L.A. Berven

275 **Downregulation of an AIM-1 kinase couples with megakaryocytic polyploidization of human hematopoietic cells.**

A. Kawasaki, I. Matsumura, J.-i. Miyagawa, S. Ezoe, H. Tanaka, Y. Terada, M. Tatsuka, T. Machii, H. Miyazaki, Y. Furukawa, and Y. Kanakura

289 **Biogenesis of porin of the outer mitochondrial membrane involves an import pathway via receptors and the general import pore of the TOM complex.**

T. Krimmer, D. Rapaport, M.T. Ryan, C. Meisinger, C.K. Kassenbrock, E. Blachly-Dyson, M. Forte, M.G. Douglas, W. Neupert, F.E. Nargang, and N. Pfanner

301 **The role of macrophages in demyelinating peripheral nervous system of mice heterozygously deficient in P0.**

S. Carenini, M. Mäurer, A. Werner, H. Blazyca, K.V. Toyka, C.D. Schmid, G. Raivich, and R. Martini

309 **The role of dynamin and its binding partners in coated pit invagination and scission.**

E. Hill, J. van der Kaay, C.P. Downes, and E. Smythe

325 **Negative regulation of Ros receptor tyrosine kinase signaling: An epithelial function of the SH2 domain protein tyrosine phosphatase SHP-1.**

H. Keilhack, M. Müller, S.-A. Böhmer, C. Frank, K.M. Weidner, W. Birchmeier, T. Ligensa, A. Berndt, H. Kosmehl, B. Günther, T. Müller, C. Birchmeier, and Frank D. Böhmer

335 **Intraarterial injection of muscle-derived $CD34^+Sca\text{-}1^+$ stem cells restores dystrophin in *mdx* mice.**

Y. Torrente, J.-P Tremblay, F. Pisati, M. Belicchi, B. Rossi, M. Sironi, F. Fortunato, M. El Fahime, M.G. D'Angelo, N.J. Caron, G. Constantin, D. Paulin, G. Scarlato, and N. Bresolin

349 **The Ndc80p complex from *Saccharomyces cerevisiae* contains conserved centromere components and has a function in chromosome segregation.**

P.A. Wigge and J.V. Kilmartin

361 **Convergence of $\alpha_v\beta_3$ integrin– and macrophage colony stimulating factor–mediated signals on phospholipase Cγ in prefusion osteoclasts.**

I. Nakamura, L. Lipfert, G.A. Rodan, and L.T. Duong

375 **A novel chromatin protein, distantly related to histone H2A, is largely excluded from the inactive X chromosome.**

B.P. Chadwick and H.F. Willard

385 **The positioning and dynamics of origins of replication in the budding yeast nucleus.**

P. Heun, T. Laroche, M.K. Raghuraman, and S.M. Gasser

401 **Mmm1p, a mitochondrial outer membrane protein, is connected to mitochondrial DNA (mtDNA) nucleoids and required for mtDNA stability.**

A.E. Aiken Hobbs, M. Srinivasan, J.M. McCaffery, and R.E. Jensen

Report

411 **Gradient of increasing affinity of importin β for nucleoporins along the pathway of nuclear import.**

I. Ben-Efraim and L. Gerace

419 ADDITIONS AND CORRECTIONS

421 ADDITIONS AND CORRECTIONS

423 ADDITIONS AND CORRECTIONS

Volume 152, Number 3, February 5, 2001

In Brief

Stem cells and mosaic diseases.
A novel Sec1 gene in cytokinesis.
How the extracellular matrix regulates anoikis.
Dystroglycans in acetylcholine receptor clustering.
Hsc70 regulates clathrin-coated vesicle disassembly.

A.W. Dove

Regular Articles

425 **Chromosome movement in mitosis requires microtubule anchorage at spindle poles.**

M.B. Gordon, L. Howard, and D.A. Compton

435 **The dystroglycan complex is necessary for stabilization of acetylcholine receptor clusters at neuromuscular junctions and formation of the synaptic basement membrane.**

C. Jacobson, P.D. Côté, S.G. Rossi, R.L. Rotundo, and S. Carbonetto

451 **A novel function of *Saccharomyces cerevisiae CDC5* in cytokinesis.**

S. Song and K.S. Lee

471 **A novel neural Wiskott-Aldrich syndrome protein (N-WASP) binding protein, WISH, induces Arp2/3 complex activation independent of Cdc42.**

M. Fukuoka, S. Suetsugu, H. Miki, K. Fukami, T. Endo, and T. Takenawa

483 **DIABLO promotes apoptosis by removing MIHA/XIAP from processed caspase 9.**

P.G. Ekert, J. Silke, C.J. Hawkins, A.M. Verhagen, and D.L. Vaux

491 **Apical membrane localization of the adenomatous polyposis coli tumor suppressor protein and subcellular distribution of the β-catenin destruction complex in polarized epithelial cells.**

A. Reinacher-Schick and B.M. Gumbiner

503 **Erv41p and Erv46p: New components of COPII vesicles involved in transport between the ER and Golgi complex.**

S. Otte, W.J. Belden, M. Heidtman, J. Liu, O.N. Jensen, and C. Barlowe

519 **Two distinct Vps34 phosphatidylinositol 3–kinase complexes function in autophagy and carboxypeptidase Y sorting in *Saccharomyces cerevisiae.***

A. Kihara, T. Noda, N. Ishihara, and Y. Ohsumi

531 **The cytokinesis gene *KEULE* encodes a Sec1 protein that binds the syntaxin KNOLLE.**

F.F. Assaad, Y. Huet, U. Mayer, and G. Jürgens

545 **PITX2 regulates procollagen lysyl hydroxylase (PLOD) gene expression: Implications for the pathology of Rieger syndrome.**

T.A. Hjalt, B.A. Amendt, and J.C. Murray

553 **Functional differences in yeast protein disulfide isomerases.**

P. Nørgaard, V. Westphal, C. Tachibana, L. Alsøe, B. Holst, and J.R. Winther

563 **Identification and characterization of an escorter for two secretory adhesins in *Toxoplasma gondii.***

M. Reiss, N. Viebig, S. Brecht, M.-N. Fourmaux, M. Soete, M. Di Cristina, J.F. Dubremetz, and D. Soldati

579 **Cdc42Hs facilitates cytoskeletal reorganization and neurite outgrowth by localizing the 58-kD insulin receptor substrate to filamentous actin.**

S. Govind, R. Kozma, C. Monfries, L. Lim, and S. Ahmed

595 **Distribution and function of AP-1 clathrin adaptor complexes in polarized epithelial cells.**

H. Fölsch, M. Pypaert, P. Schu, and I. Mellman

607 **Dominant-interfering hsc70 mutants disrupt multiple stages of the clathrin-coated vesicle cycle in vivo.**

S.L. Newmyer and S.L. Schmid

621 **Tortoise, a novel mitochondrial protein, is required for directional responses of *Dictyostelium* in chemotactic gradients.**

S. van Es, D. Wessels, D.R. Soll, J. Borleis, and P.N. Devreotes

633 **Matrix attachment regulates Fas-induced apoptosis in endothelial cells: A role for c-Flip and implications for anoikis.**

F. Aoudjit and K. Vuori

Reports

645 **Focal activation of a mutant allele defines the role of stem cells in mosaic skin disorders.**

M.J. Arin, M.A. Longley, X.-J. Wang, and D.R. Roop

651 **An inducible mouse model for epidermolysis bullosa simplex: Implications for gene therapy.**

T. Cao, M.A. Longley, X.-J. Wang, and D.R. Roop

Volume 152, Number 4, February 19, 2001

In Brief

Tyrosine phosphorylation of IκBα.
Visualizing autophagy.
MMPs as inhibitors of adipocyte differentiation.
Degradation and aggregation of an amyloidogenic protein.
The ins and outs of karyopherins.

A.W. Dove

Comments

F17 **Where Notch and Wnt signaling meet: The presenilin hub.**

B. De Strooper and W. Annaert

F21 **Of yeast, mice, and men: Rab proteins and organelle transport.**

S.W. Deacon and V.I. Gelfand

Regular Articles

657 **Dissection of autophagosome formation using Apg5-deficient mouse embryonic stem cells.**

N. Mizushima, A. Yamamoto, M. Hatano, Y. Kobayashi, Y. Kabeya, K. Suzuki, T. Tokuhisa, Y. Ohsumi, and T. Yoshimori

669 ***Drosophila* Aurora B kinase is required for histone H3 phosphorylation and condensin recruitment during chromosome condensation and to organize the central spindle during cytokinesis.**

R. Giet and D.M. Glover

683 **Connection of the mitochondrial outer and inner membranes by Fzo1 is critical for organellar fusion.**

S. Fritz, D. Rapaport, E. Klanner, W. Neupert, and B. Westermann

693 **Stromelysin-1 regulates adipogenesis during mammary gland involution.**

C.M. Alexander, S. Selvarajan, J. Mudgett, and Z. Werb

705 **Hsp70 and antifibrillogenic peptides promote degradation and inhibit intracellular aggregation of amyloidogenic light chains.**

J.L. Dul, D.P. Davis, E.K. Williamson, F.J. Stevens, and Y. Argon

717 **A specific role of phosphatidylinositol 3–kinase γ: A regulation of autonomic Ca^{2+} oscillations in cardiac cells.**

C. Bony, S. Roche, U. Shuichi, T. Sasaki, M.A. Crackower, J. Penninger, H. Mano, and M. Pucéat

729 **The karyopherin Kap142p/Msn5p mediates nuclear import and nuclear export of different cargo proteins.**

K. Yoshida and G. Blobel

741 **Plasminogen activator inhibitor 1 functions as a urokinase response modifier at the level of cell signaling and thereby promotes MCF-7 cell growth.**

D.J. Webb, K.S. Thomas, and S.L. Gonias

753 **Activation of nuclear factor κB and *bcl-x* survival gene expression by nerve growth factor requires tyrosine phosphorylation of IκBα.**

N.T. Bui, A. Livolsi, J.-F. Peyron, and J.H.M. Prehn

765 **A novel 14-kilodalton protein interacts with the mitogen-activated protein kinase scaffold MP1 on a late endosomal/lysosomal compartment.**

W. Wunderlich, I. Fialka, D. Teis, A. Alpi, A. Pfeifer, R.G. Parton, F. Lottspeich, and L.A. Huber

777 **The plasminogen activator inhibitor PAI-1 controls in vivo tumor vascularization by interaction with proteases, not vitronectin: Implications for antiangiogenic strategies.**

K. Bajou, V. Masson, R.D. Gerard, P.M. Schmitt, V. Albert, M. Praus, L.R. Lund, T.L. Frandsen, N. Brunner, K. Dano, N.E. Fusenig, U. Weidle, G. Carmeliet, D. Loskutoff, D. Collen, P. Carmeliet, J.M. Foidart, and A. Noël

785 **Presenilin 1 negatively regulates β-catenin/T cell factor/lymphoid enhancer factor-1 signaling independently of β-amyloid precursor protein and notch processing.**

S. Soriano, D.E. Kang, M. Fu, R. Pestell, N. Chevallier, H. Zheng, and E.H. Koo

795 **Rab27a regulates the peripheral distribution of melanosomes in melanocytes.**

A.N. Hume, L.M. Collinson, A. Rapak, A.Q. Gomes, C.R. Hopkins, and M.C. Seabra

809 **Distinct protein sorting and localization to premelanosomes, melanosomes, and lysosomes in pigmented melanocytic cells.**

G. Raposo, D. Tenza, D.M. Murphy, J.F. Berson, and M.S. Marks

825 **Rab27a is required for regulated secretion in cytotoxic T lymphocytes.**

J.C. Stinchcombe, D.C. Barral, E.H. Mules, S. Booth, A.N. Hume, L.M. Machesky, M.C. Seabra, and G.M. Griffiths

835 **Defective granule exocytosis in Rab27a-deficient lymphocytes from *Ashen* mice.**

E.K. Haddad, X. Wu, J.A. Hammer III, and P.A. Henkart

Reports

843 **Rab27a: A key to melanosome transport in human melanocytes.**

P. Bahadoran, E. Aberdam, F. Mantoux, R. Buscà, K. Bille, N. Yalman, G. de Saint-Basile, R. Casaroli-Marano, J.-P. Ortonne, and R. Ballotti

851 **An essential role for the substrate-binding region of Hsp40s in *Saccharomyces cerevisiae.***

J.L. Johnson and E.A. Craig

Volume 152, Number 5, March 5, 2001

In Brief

Oscillating actin.
A cargo for kinesin.
Microtubules in the end-zone.
Linking without signaling.
A stretchy structure.
W.A. Wells

Comments

F25 **Kinesin delivers: Identifying receptors for motor proteins.**
P.J. Hollenbeck

F29 **Caveolin, cholesterol, and lipid droplets?**
G. van Meer

Regular Articles

857 **α_4-Integrin mediates neutrophil-induced free radical injury to cardiac myocytes.**
B.Y. Poon, C.A. Ward, C.B. Cooper, W.R. Giles, A.R. Burns, and P. Kubes

867 **Phosphatidylinositol 4,5-bisphosphate induces actin stress-fiber formation and inhibits membrane ruffling in CV1 cells.**
M. Yamamoto, D.H. Hilgemann, S. Feng, H. Bito, H. Ishihara, Y. Shibasaki, and H.L. Yin

877 **A novel interaction of the Golgi complex with the vimentin intermediate filament cytoskeleton.**
Y.-s. Gao and E. Sztul

895 **Cofactor requirements for nuclear export of Rev response element (RRE)– and constitutive transport element (CTE)–containing retroviral RNAs: An unexpected role for actin.**
W. Hofmann, B. Reichart, A. Ewald, E. Müller, I. Schmitt, R.H. Stauber, F. Lottspeich, B.M. Jockusch, U. Scheer, J. Hauber, and M.-C. Dabauvalle

911 **Autocrine stimulation of human pancreatic duct–like development by soluble isoforms of epimorphin in vitro.**
L. Lehnert, M.M. Lerch, Y. Hirai, M.-L. Kruse, W. Schmiegel, and H. Kalthoff

923 **The Golgi-associated Hook3 protein is a member of a novel family of microtubule-binding proteins.**
J.H. Walenta, A.J. Didier, X. Liu, and H. Krämer

935 **Rer1p, a retrieval receptor for endoplasmic reticulum membrane proteins, is dynamically localized to the Golgi apparatus by coatomer.**
K. Sato, M. Sato, and A. Nakano

945 **The localization of human cyclins B1 and B2 determines CDK1 substrate specificity and neither enzyme requires MEK to disassemble the Golgi apparatus.**
V.M. Draviam, S. Orrechia, M. Lowe, R. Pardi, and J. Pines

959 **Cargo of kinesin identified as JIP scaffolding proteins and associated signaling molecules.**
K.J. Verhey, D. Meyer, R. Deehan, J. Blenis, B.J. Schnapp, T.A. Rapoport, and B. Margolis

971 **Regulation of CDC42 GTPase by proline-rich tyrosine kinase 2 interacting with PSGAP, a novel pleckstrin homology and Src homology 3 domain containing rhoGAP protein.**
X.-R. Ren, Q.-S. Du, Y.-Z. Huang, S.-Z. Ao, L. Mei, and W.-C. Xiong

985 **Evidence for a replication function of FFA-1, the *Xenopus* orthologue of Werner syndrome protein.**
C.-Y. Chen, J. Graham, and H. Yan

997 **Regulation of antioxidant metabolism by translation initiation factor 2α.**
S. Tan, N. Somia, P. Maher, and D. Schubert

1007 ***Drosophila* integrin-linked kinase is required at sites of integrin adhesion to link the cytoskeleton to the plasma membrane.**
C.G. Zervas, S.L. Gregory, and N.H. Brown

1019 **Rop GTPase–dependent dynamics of tip-localized F-actin controls tip growth in pollen tubes.**
Y. Fu, G. Wu, and Z. Yang

1033 **Protein kinase C activation promotes microtubule advance in neuronal growth cones by increasing average microtubule growth lifetimes.**
N. Kabir, A.W. Schaefer, A. Nakhost, W.S. Sossin, and P. Forscher

1045 **The supramolecular organization of fibrillin-rich microfibrils.**
C. Baldock, A.J. Koster, U. Ziese, M.J. Rock, M.J. Sherratt, K.E. Kadler, C.A. Shuttleworth, and C.M. Kielty

1057 **A caveolin dominant negative mutant associates with lipid bodies and induces intracellular cholesterol imbalance.**
A. Pol, R. Luetterforst, M. Lindsay, S. Heino, E. Ikonen, and R.G. Parton

1071 **Accumulation of caveolin in the endoplasmic reticulum redirects the protein to lipid storage droplets.**
A.G. Ostermeyer, J.M. Paci, Y. Zeng, D.M. Lublin, S. Munro, and D.A. Brown

1079 **Caveolin-2 is targeted to lipid droplets, a new "membrane domain" in the cell.**
T. Fujimoto, H. Kogo, K. Ishiguro, K. Tauchi, and R. Nomura

1087 **Mammalian Sprouty-1 and -2 are membrane-anchored phosphoprotein inhibitors of growth factor signaling in endothelial cells.**
M.-A. Impagnatiello, S. Weitzer, G. Gannon, A. Compagni, M. Cotten, and G. Christofori

Reports

1099 **The transcription coactivator CBP is a dynamic component of the promyelocytic leukemia nuclear body.**
F.-M. Boisvert, M.J. Kruhlak, A.K. Box, M.J. Hendzel, and D.P. Bazett-Jones

1107 **Deletion of murine *SMN* exon 7 directed to skeletal muscle leads to severe muscular dystrophy.**
C. Cifuentes-Diaz, T. Frugier, F.D. Tiziano, E. Lacène, N. Roblot, V. Joshi, M.H. Moreau, and J. Melki

1115 **Equine herpesvirus protein E10 induces membrane recruitment and phosphorylation of its cellular homologue, bcl-10.**
M. Thome, O. Gaide, O. Micheau, F. Martinon, D. Bonnet, M. Gonzalez, and J. Tschopp

Volume 152, Number 6, March 19, 2001

In Brief

Angiostatin antagonizes angiomotin.
Endostatin antagonizes itself, trimerized.
Integrins to the rescue.
HMG1 and atherosclerosis.
W.A. Wells

Comment

F35 **Hold that line: Angiomotin regulates endothelial cell motility.**
B.R. Zetter

Regular Articles

1123 ***UGO1* encodes an outer membrane protein required for mitochondrial fusion.**
H. Sesaki and R.E. Jensen

1135 **Diacylglycerol kinase ζ regulates Ras activation by a novel mechanism.**
M.K. Topham and S.M. Prescott

1145 **Rac mediates cytoskeletal rearrangements and increased cell motility induced by urokinase-type plasminogen activator receptor binding to vitronectin.**
L. Kjøller and A. Hall

1159 **An acidic motif retains vesicular monoamine transporter 2 on large dense core vesicles.**
C.L. Waites, A. Mehta, P.K. Tan, G. Thomas, R.H. Edwards, and D.E. Krantz

1169 **A role for syndecan-1 in coupling fascin spike formation by thrombospondin-1.**
J.C. Adams, N. Kureishy, and A.L. Taylor

1183 **Atypical protein kinase C is involved in the evolutionarily conserved PAR protein complex and plays a critical role in establishing epithelia-specific junctional structures.**
A. Suzuki, T. Yamanaka, T. Hirose, N. Manabe, K. Mizuno, M. Shimizu, K. Akimoto, Y. Izumi, T. Ohnishi, and S. Ohno

1197 **The high mobility group (HMG) boxes of the nuclear protein HMG1 induce chemotaxis and cytoskeleton reorganization in rat smooth muscle cells.**
B. Degryse, T. Bonaldi, P. Scaffidi, S. Müller, M. Resnati, F. Sanvito, G. Arrigoni, and M.E. Bianchi

1207 **Enhanced expression of the α7β1 integrin reduces muscular dystrophy and restores viability in dystrophic mice.**
D.J. Burkin, G.O. Wallace, K.J. Nicol, D.J. Kaufman, and S.J. Kaufman

1219 **The NC1/endostatin domain of *Caenorhabditis elegans* type XVIII collagen affects cell migration and axon guidance.**
B.D. Ackley, J.R. Crew, H. Elamaa, T. Pihlajaniemi, C.J. Kuo, and J.M. Kramer

1233 **Oligomerization-dependent regulation of motility and morphogenesis by the collagen XVIII NC1/endostatin domain.**
C.J. Kuo, K.R. LaMontagne, Jr., G. Garcia-Cardeña, B.D. Ackley, D. Kalman, S. Park, R. Christofferson, J. Kamihara, Y.-H. Ding, K.-M. Lo, S. Gillies, J. Folkman, R.C. Mulligan, and K. Javaherian

1247 **Angiomotin: An angiostatin binding protein that regulates endothelial cell migration and tube formation.**
B. Troyanovsky, T. Levchenko, G. Månsson, O. Matvijenko, and L. Holmgren

1255 **Budding yeast chromosome structure and dynamics during mitosis.**
C.G. Pearson, P.S. Maddox, E.D. Salmon, and K. Bloom

1267 **Cyclin E uses Cdc6 as a chromatin-associated receptor required for DNA replication.**
L. Furstenthal, B.K. Kaiser, C. Swanson, and P.K. Jackson

1279 **Internal modification of U2 small nuclear (sn)RNA occurs in nucleoli of *Xenopus* oocytes.**
Y.-T. Yu, M.-D. Shu, A. Narayanan, R.M. Terns, M.P. Terns, and J.A. Steitz

1289 **Glial growth factor/neuregulin inhibits Schwann cell myelination and induces demyelination.**
G. Zanazzi, S. Einheber, R. Westreich, M.-J. Hannocks, D. Bedell-Hogan, M.A. Marchionni, and J.L. Salzer

Reports

1301 **Oncogenic Ras-induced proliferation requires autocrine fibroblast growth factor 2 signaling in skeletal muscle cells.**
Y.V. Fedorov, R.S. Rosenthal, and B.B. Olwin

1307 **Importin β–mediated nuclear import of fibroblast growth factor receptor: Role in cell proliferation.**
J.F. Reilly and P.A. Maher

1313 **The *Caenorhabditis elegans unc-78* gene encodes a homologue of actin-interacting protein 1 required for organized assembly of muscle actin filaments.**
S. Ono